

# Mapping of the unoccupied states and relevant bosonic modes via the time dependent momentum distribution

A. F. Kemper,<sup>1,2,\*</sup> M. Sentef,<sup>2</sup> B. Moritz,<sup>3,2,4</sup> C. C. Kao,<sup>5</sup> Z. X. Shen,<sup>2,6</sup> J. K. Freericks,<sup>7</sup> and T. P. Devereaux<sup>2,6</sup>

<sup>1</sup>*Computational Research Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA*

<sup>2</sup>*Stanford Institute for Materials and Energy Sciences,*

*SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA*

<sup>3</sup>*Department of Physics, Northern Illinois University, DeKalb, IL 60115, USA*

<sup>4</sup>*Department of Physics and Astrophysics, University of North Dakota, Grand Forks, ND 58202, USA*

<sup>5</sup>*Stanford Synchrotron Radiation Lightsource, SLAC National Accelerator Laboratory, Menlo Park, California 94025, USA*

<sup>6</sup>*Geballe Laboratory for Advanced Materials, Stanford University, Stanford, California 94305, USA*

<sup>7</sup>*Department of Physics, Georgetown University, Washington, DC 20057, USA*

The unoccupied states of complex materials are difficult to measure, yet play a key role in determining their emergent properties. We propose a novel technique that can measure the unoccupied states, time-resolved Compton scattering, which measures the time-dependent momentum distribution (TDMD). Using a non-equilibrium Keldysh formalism, we study the TDMD for electrons coupled to a lattice in a pump-probe setup. We find a direct relation between temporal oscillations in the TDMD and the underlying unoccupied states, suggesting that both can be measured by time-resolved Compton scattering.

PACS numbers: 78.47.je, 79.60.-i, 63.20.kd

In understanding the emergent properties of complex materials, it is insufficient to limit ones' study solely to the occupied electronic states. The unoccupied states play an important role in determining, for example, the nature of the gap in charge- and spin-density waves, the absorption properties of semiconductors, or the magnetic properties of Mott systems. For example, without knowledge of the unoccupied states, it is difficult to measure both the full width as well as the  $\vec{q}$  of a gap in the electronic spectrum. Unfortunately, the most direct experimental measurements of momentum-resolved states in quantum materials, angle-resolved photoemission spectroscopy (ARPES), only determines the occupied states. This leaves the alternative methods of inverse photoemission spectroscopy (IPES), which lacks the signal strength to measure with sufficient resolution, or indirectly inferring the unoccupied dispersion from two-particle quantities such as optical spectroscopy. Pump-probe spectroscopy, and in particular time-resolved (tr-) ARPES, can measure part of the unoccupied states, but is limited by the inherent competition between temporal and energy resolution, and cannot map out the entire Brillouin zone in most cases.

Here we consider a different quantity, where one sacrifices the energy information in favor of high time resolution, while maintaining high momentum resolution. We propose to measure the (gauge-invariant) time-dependent momentum distribution (TDMD) of quasiparticles, defined by  $\langle c_{\mathbf{k}}^{\dagger}(t)c_{\mathbf{k}}(t) \rangle \equiv n_{\mathbf{k}}(t)$ , which is similar to the so-called Wigner distribution. It contains a rich amount of information about the behavior of pumped quasiparticles while avoiding some of the resolution-related complications of other measurements.

The TDMD is commonly measured via time-of-flight

absorption images in cold atomic gases, where it was recently used to map out the Fermi surface of a gas of  $^{40}\text{K}$  atoms.[1] In the context of condensed matter systems, we propose two new types of measurements to access the TDMD. First, we propose to extend the technique of Compton scattering for solids into the time domain[2]. Compton scattering has a long history as a probe of the electron momentum distribution for solids in equilibrium [3–5]. With the advent of x-ray free-electron laser sources, the photon energy of the ultrashort x-ray probe pulses has been extended into the hard x-ray regime, where the fractional Compton cross-section becomes appreciable. The planned upgrade of the Linac Coherent Light Source at SLAC National Accelerator Laboratory will extend the energy of the x-ray free electron laser to beyond 20 keV. There is also significant progress in crystal optics at these energies, such that time-resolved high resolution Compton scattering with momentum resolution of a few percent of the Brillouin zone will be feasible. Additionally, recent developments in photonics suggest the future availability of ultrashort pulses ranging from x-rays to gamma rays[6]. Time-resolved Compton scattering has the advantage of using high-momentum photons, which can readily access the Brillouin zone edges in all directions. Secondly, one can access the TDMD with tr-ARPES by using similar time delays and probe pulse widths and integrating the tr-ARPES signal over energy.

We illustrate the use of the TDMD by demonstrating how it can be used to map out the unoccupied dispersion in a pump-probe setup on a model system with electron-lattice interactions appropriate to a large class of complex materials. The TDMD exhibits two ubiquitous phenomena in time-resolved experiments; both are due to electron-phonon interactions. First, the phonons




FIG. 1. a) (Color online) Equilibrium TDMD  $n_{\mathbf{k}}(t)$  far before the pump arrives. The gray area indicates where quasiparticle states are occupied. The Fermi surface is sharply defined for the parameters presented ( $g^2 = \sqrt{0.02}V_{nn}$ ,  $T = 0.04V_{nn}$ ,  $\Omega = 0.4V_{nn}$ ). b) Change in the momentum distribution from equilibrium  $\delta n_{\mathbf{k}}(t)$  after a pump at  $t \approx 4.8\tau_p$ . The arrow indicates the direction of the applied electric field.

dissipate the energy delivered to the electronic system by the pump. Second, after pumping, oscillations related to the phonon frequencies (and independent of the initial pump parameters) are commonly observed in the resulting spectra measured by the probe [7–10]. We will show that both of these features are clearly seen in the TDMD.

We study a two-dimensional system of electrons coupled to a bath of non-dispersive phonons with frequency  $\Omega$  via a constant coupling  $g$  (known as the Holstein model), [11].

$$\mathcal{H} = \sum_{\mathbf{k}\sigma} (\epsilon(\mathbf{k}) - \mu) c_{\mathbf{k},\sigma}^\dagger c_{\mathbf{k},\sigma} + \sum_{\mathbf{q}} \Omega b_{\mathbf{q}}^\dagger b_{\mathbf{q}} + g \sum_{\mathbf{k}\mathbf{q}\sigma} c_{\mathbf{k}+\mathbf{q},\sigma}^\dagger c_{\mathbf{k},\sigma} (b_{\mathbf{q}} + b_{-\mathbf{q}}^\dagger) \quad (1)$$

where  $c_{\mathbf{k}}$  ( $b_{\mathbf{q}}$ ) annihilates an electron (phonon) of momentum  $\mathbf{k}$  ( $\mathbf{q}$ ).

While the TDMD is useful independent of the underlying model, here we use a band structure motivated by the transition-metal oxides: a 2D tight-binding  $\epsilon(\mathbf{k})$  with (next) nearest-neighbor hoppings  $V_{nn}$  and  $V_{nnn} = 0.3V_{nn}$ , and  $\mu = -1.02V_{nn}$ . We use the conventions  $\hbar = e = c = 1$ . Below, we will report timescales in units of a characteristic phonon timescale  $\tau_p = 1/\Omega$ . For example, a phonon with an energy of  $\Omega = 10$  meV has  $\tau_p \approx 0.4$  ps. Similarly, there is a characteristic electron timescale  $\tau_e = 1/V_{nn} \approx 16$  fs for  $V_{nn} = 250$  meV. Although in this work we will explicitly examine higher energy phonons for numerical stability reasons, the results below occur

due to the relevant energy scales in the problem, and will simply be rescaled to lower phonon energies.

To describe the non-equilibrium pump-probe process, we propagate the system on the Kadanoff-Baym-Keldysh (Keldysh) contour [12], which has been described in detail elsewhere [13–15]. The non-equilibrium Keldysh formalism has been successfully used to describe the time evolution of correlated electrons within dynamical mean-field theory [16–19]. The electric field is included through the standard Peierls' substitution  $\mathbf{k} \rightarrow \mathbf{k} - \mathbf{A}(t)$ , where the vector potential  $\mathbf{A}$  is related to the applied electric field  $\mathbf{F}$  via  $-\partial \mathbf{A}(t)/\partial t = \mathbf{F}$  and we work in the Hamiltonian gauge.

Within the Migdal limit, we consider the renormalization of the electron Green's function by a single phonon emission/absorption and ignore the phonon self-energy (as is done in conventional Migdal-Eliashberg theory). In this limit, we solve the Dyson equation

$$G_{\mathbf{k}}(t, t') = G_{\mathbf{k}}^0(t, t') + \int_{\mathcal{C}} dt_1 dt_2 G_{\mathbf{k}}^0(t, t_1) \Sigma(t_1, t_2) G_{\mathbf{k}}(t_2, t')$$

with the self-energy  $\Sigma(t, t') = ig^2 \sum_{\mathbf{k}} D^0(t, t') G_{\mathbf{k}}^0(t, t')$ , where  $G_{\mathbf{k}}^0(t, t')$  and  $D^0(t, t')$  are the non-interacting electron and phonon Green's functions, respectively [13]. The Dyson equation can be solved by recasting it as a matrix equation [15] or by decomposing into Volterra-type equations through the Langreth rules [20], which is the approach used here.

Figure 1 shows the gauge-invariant TDMD  $n_{\mathbf{k}}(t) =$

$iG_{\mathbf{k}+\vec{A}(t)}^<(t, t' \rightarrow t)$ [13] for  $g = \sqrt{0.02}V_{nn}$ ,  $\Omega = 0.4V_{nn}$  and at a temperature  $T = 0.04V_{nn}$ . Fig. 1a) shows  $n_{\mathbf{k}}(t)$  at times far before the pump, where the system is fully described by the equilibrium problem. To clearly show the changes for the small pump fluences considered ( $F_{\max} \leq 4V_{nn}$ ) Fig. 1b) shows the change in the TDMD from equilibrium:  $\delta n_{\mathbf{k}}(t) \equiv n_{\mathbf{k}}(t) - n_{\mathbf{k}}(t \rightarrow -\infty)$ .

We apply a pulse centered at time  $t = 0$  in the (11) (diagonal) direction of the form  $A(t) = (F_{\max}/\omega_A) \exp(-t^2/(2\sigma^2)) \sin(\omega_A t)$ , with a maximum field strength of  $F_{\max} = 4V_{nn}$ , a frequency  $\omega_A = 2V_{nn}$  and  $\sigma \approx 0.4\tau_p$ . At the time shown in Fig. 1 b), the pump has passed, and the system is relaxing towards equilibrium. The  $C_4$  symmetry of  $n_{\mathbf{k}}(t)$  in equilibrium is broken by the field (which points in the (11) direction as shown by the black arrow), and this is reflected in the transient change. In fact, this can be used to emphasize particular regions of interest in the Brillouin zone. By rotating the field, the regions exhibiting the largest change in  $n_{\mathbf{k}}(t)$  will shift with the field direction, providing access to new regions by emphasizing the experimental signal near different momenta.




FIG. 2. a) (Color online) Change in the TDMD  $n_{\mathbf{k}}(t)$  from the equilibrium value (at  $t = 0$ ) for the points in the Brillouin zone indicated in b). Symbols are calculated data points (for clarity, not all are shown), solid lines are fits to an exponential decay using a fixed relaxation rate  $1/\tau_{\mathbf{k}} = -2\text{Im } \Sigma(\omega = \epsilon_{\mathbf{k}})$ . c) Real and imaginary parts of the equilibrium self-energy  $\Sigma(\omega)$  for the parameters in a) ( $g = \sqrt{0.02}V_{nn}$ ,  $T = 0.04V_{nn}$ ,  $\Omega = 0.4V_{nn}$ ).

First, we consider the anisotropic redistribution of quasiparticles as a function of time. In Figure 2a), we show  $\delta n_{\mathbf{k}}(t)$  at the momenta indicated in the inset; the blue (black) momenta have the same energy  $\epsilon_{\mathbf{k}}$  but due to the  $C_4$  symmetry breaking have different time traces. From the figure, one can clearly see that there is a difference in the overall magnitude of the change at the

equivalent momenta (with the same color). The solid lines are fits to the amplitude of a simple decaying exponential, where we have taken the decay constant at a momentum  $\mathbf{k}$  from the on-shell *equilibrium* self-energy  $1/\tau_{\mathbf{k}}(\omega) = -2\text{Im } \Sigma(\omega = \epsilon_{\mathbf{k}})$  shown in Fig. 2b), as suggested by a recent study[21]. The remarkable agreement with the calculated data points confirms that it is indeed the equilibrium self-energy that controls the decay rate. Since the equilibrium self-energy is  $C_4$  symmetric, the anisotropic redistribution in  $\delta n_{\mathbf{k}}(t)$  is caused by the  $C_4$  symmetry breaking of the applied pump field.




FIG. 3. a) (Color online) Oscillatory part of  $\delta n_{\mathbf{k}}(t)$  for  $\mathbf{k}$  along (11) at the energies  $\epsilon_{\mathbf{k}}$  indicated (offset for clarity). b) Fourier transform of a). Vertical lines indicate the frequencies corresponding to the strong oscillations on the unoccupied side,  $\omega = \epsilon_{\mathbf{k}} + \Omega$  and  $\omega = \epsilon_{\mathbf{k}} - W_- + \Omega$ . In both a) and b) the oscillations on the unoccupied side close to the Fermi level (blue curves) have been scaled down by 4 for visibility. (see text for details).

A ubiquitous feature observed in pump-probe experiments is oscillations[7–10] of the time traces dominated by a few characteristic frequencies. These are clearly observed as rings in  $\delta n_{\mathbf{k}}(t)$  shown in Fig. 1, as well as oscillations in the time traces in Fig. 2. We further emphasize these in Fig. 3, where we have subtracted the decaying exponential from several points along the Brillouin zone diagonal in Fig. 1. From Fig. 3a), one can observe two

main features. First, the oscillation frequencies in the time traces depend strongly on momentum. Second, although there is only a single phonon frequency in our model, several oscillation frequencies can be observed. Figure 3b) shows the Fourier transform power spectrum of the oscillations in Fig. 3a). Each curve has readily visible maxima, in addition to some smaller structure across the frequency spectrum. For each energy (momentum)  $\epsilon_{\mathbf{k}}$ , the vertical lines indicate the energies  $\omega = \epsilon_{\mathbf{k}} + \Omega$  and  $\omega = \epsilon_{\mathbf{k}} - W_- + \Omega$ , where  $W_-$  is the energy at the bottom of the band (at  $\mathbf{k} = \Gamma$ ).

The temporal dynamics of the Holstein model leads to this rich physics. The decay and the oscillations observed are momentum (energy) dependent, and their amplitudes depend on the electron-phonon coupling, as well as the applied field. Describing the full temporal dynamics of  $n_{\mathbf{k}}(t)$  is complicated, yet we can gain some insight by focusing on the times where the pump is off, while the system is still displaced from its equilibrium configuration. The subsequent relaxation occurs from the new configuration through the equations of motion. In equilibrium, the TDMD is balanced by equal rates of in- and out-scattering; once the pump breaks this symmetry, the system relaxes back to the equilibrium according to the dynamics contained within the self-energy and Green's functions. In addition to the relaxation, the TDMD oscillates at the band energy shifted by the characteristic energies in the problem (the phonon frequency and bandwidth) at which the self-energy is largest. This is shown analytically at  $T = 0$  in the supplemental information; the full calculation requires the numerical methods used here.

At low temperatures, the real part of the self-energy  $\text{Re } \Sigma$  is large and peaked near four frequencies,  $\omega = \pm\Omega$  and  $\omega = W_{\pm} \pm \Omega$ , where  $W_{\pm}$  is the upper (lower) band edge (see Fig. 2). The resulting  $n_{\mathbf{k}}(t)$  oscillates at the band energy  $\epsilon_{\mathbf{k}}$  shifted by those frequencies. In principle, all four frequencies should be observable in the power spectrum of Fig. 3; however, two frequencies (with the positive sign above) correspond to phonon absorption, which is small at low temperatures, and thus the corresponding peaks in the power spectrum are reduced significantly. The two strong peaks expected for the unoccupied side are indicated in Fig. 3 by vertical lines, which agree well with the observed frequencies, up to a small shift in energy due to  $\text{Re } \Sigma$ . An interesting consequence of the expected oscillation frequencies is that if one considers a measurement near the Fermi level, the oscillations will appear to be strongest at just the phonon frequency (as observed in tr-ARPES[7–10]).

The simple dependence of the oscillation frequencies on the underlying band structure suggests a novel method to measure the electron dispersion by looking at the oscillations in the TDMD. The oscillations are most clearly visible in the unoccupied region of the Brillouin zone, which is exactly the region that traditional methods for

measuring the dispersion have difficulty accessing. By tracking the oscillation frequencies as a function of momentum one can directly map out the dispersion. In Fig. 4, we plot the Fourier transform power spectrum in the unoccupied portion of the Brillouin zone along the zone diagonal and zone face. This shows most clearly the two strong peaks in the Fourier transform associated with phonon emission, although a weaker line corresponding to phonon absorption is also visible below the two main lines. All three lines, up to the constant shifts from the real part of the self-energy described above, follow the unoccupied dispersion. Thus, from the maxima in the power spectrum the unoccupied dispersion can be directly measured.

The above analysis, applied to theoretical results, can be applied to experimental results equally well, although one must take care to include the  $\alpha^2 F(\Omega)$  for the material under study. Time-resolved Compton scattering also requires the introduction of the scattering matrix elements, similar to what has been addressed in equilibrium Compton scattering[3]. Additionally, Compton scattering measures a single momentum direction, averaging over the directions not selected by the field. This allows for selectivity in the direction of interest, focusing on the regions of the Brillouin zone that one wishes to measure. Finally, there is a distinction between the crystal momentum in the TDMD and the real momentum measured by Compton scattering. The inversion from real momentum to crystal momentum has been addressed in some detail previously.[22]

Experimentally, the width of the probe pulse limits the temporal resolution of the TDMD. Thus the Fourier transform to frequencies is limited in turn to several hundred meV around the Fermi level. However, in many systems of interest, the features one wishes to investigate lie near the Fermi level, both above and below. For example, in the high-Tc cuprates, several features are predicted to lie within this energy range.[23] In other gapped systems, without the knowledge of the spectra above the Fermi level it is impossible to correctly assign the gap magnitude. Similar effects due to spin fluctuations, phonons, and other interesting collective and bosonic features lie at low energy, and are thus accessible by this technique.

*Acknowledgments* AFK, MS, BM and TPD were supported by the U.S. Department of Energy, Basic Energy Sciences, Materials Sciences and Engineering Division under contract No. DE-AC02-76SF00515. JKF was supported by the U.S. Department of Energy, Basic Energy Sciences, Materials Sciences and Engineering Division under contract No. DE-FG02-08ER46542 and by the McDevitt bequest at Georgetown University. The collaboration was supported by the U.S. Department of Energy, Basic Energy Sciences, Materials Sciences and Engineering Division under contract Nos. DE-FG02-08ER46540 and de-sc0007091. This work was made possible by the resources of the National Energy




FIG. 4. (Color online) Fourier power spectrum of the oscillation frequencies along the  $\Gamma$ -M and M-X directions in the unoccupied Brillouin zone. The two strong peaks in the Fourier spectra follow the lines corresponding to phonon emission processes; a small tertiary line of peaks can be seen at lower frequencies corresponding to a phonon absorption process. The Fermi surface is shown in green on the base plane.

Research Scientific Computing Center which is supported by the U.S. DOE, Office of Science, under Contract No. DE-AC02-05CH11231. We gratefully acknowledge discussions with P. S. Kirchmann, J. Sobota, M. Wolf, and S. Yang.

\* afkemper@lbl.gov

- [1] T. E. Drake, Y. Sagi, R. Paudel, J. T. Stewart, J. P. Gaebler, and D. S. Jin, preprint (2012), arXiv:1204.0048.
- [2] Time-resolved Compton scattering in 1D was previously studied by R. E. Wagner, Q. Su, and R. Grobe, *Phys. Rev. A* **82**, 022719 (2010).
- [3] M. J. Cooper, *Reports on Progress in Physics* **48**, 415 (1985).
- [4] A. Bansil, B. Barbiellini, S. Kaprzyk, and P. Mijnders, *J. Phys. Chem. of Solids* **62**, 2191 (2001).
- [5] Y. Sakurai, M. Itou, B. Barbiellini, P. E. Mijnders, R. S. Markiewicz, S. Kaprzyk, J. M. Gillet, S. Wakimoto, M. Fujita, S. Basak, Y. J. Wang, W. Al-Sawai, H. Lin, A. Bansil, and K. Yamada, *Science* **332**, 698 (2011).
- [6] K. Ta Phuoc, S. Corde, C. Thauray, V. Malka, A. Tafzi, J. P. Goddet, R. C. Shah, S. Sebban, and A. Rousse, *Nature Photonics* **6**, 308 (2012).
- [7] F. Schmitt, P. S. Kirchmann, U. Bovensiepen, R. G. Moore, L. Rettig, M. Krenz, J.-H. Chu, N. Ru, L. Perfetti, D. H. Lu, M. Wolf, I. R. Fisher, and Z.-X. Shen, *Science* **321**, 1649 (2008).
- [8] L. Perfetti, P. A. Loukakos, M. Lisowski, U. Bovensiepen, M. Wolf, H. Berger, S. Biermann, and A. Georges, *New Journal of Physics* **10**, 053019 (2008).
- [9] L. Rettig, R. Cortés, S. Thirupathaiah, P. Gegenwart, H. S. Jeevan, T. Wolf, U. Bovensiepen, M. Wolf, H. A. Dürr, and J. Fink, preprint (2010), arXiv:1008.1561.
- [10] K. W. Kim, A. Pashkin, H. Schäfer, M. Beyer, M. Porer, T. Wolf, C. Bernhard, J. Demsar, R. Huber, and A. Leitenstorfer, *Nature Materials* (2012).
- [11] T. Holstein, *Ann. Phys.* **8**, 325 (1959); **8**, 343 (1959).
- [12] L. P. Kadanoff and G. Baym, *Quantum Statistical Mechanics* (W. A. Benjamin, Inc., NY, 1962); L. V. Keldysh, *Zh. Eksp. Teor. Fiz.* **47**, 1515 (1964).
- [13] For a complete description of the non-interacting Green's functions see e.g. G. Mahan, *Many-Particle Physics* (Plenum Press, NY, 1990).
- [14] A. P. Jauho and J. W. Wilkins, *Phys. Rev. B* **29**, 1919 (1984); J. H. Davies and J. W. Wilkins, *Phys. Rev. B* **38**, 1667 (1988).
- [15] V. Turkowski and J. K. Freericks, *Phys. Rev. B* **71**, 085104 (2005); J. K. Freericks, *Phys. Rev. B* **77**, 075109 (2008).
- [16] M. Eckstein and M. Kollar, *Phys. Rev. Lett.* **100**, 120404 (2008); M. Eckstein, A. Hackl, S. Kehrein, M. Kollar, M. Moockel, P. Werner, and F. A. Wolf, *Eur. Phys. J.: S.T.* **180**, 217 (2009); M. Eckstein, M. Kollar, and P. Werner, *Phys. Rev. Lett.* **103**, 056403 (2009).
- [17] B. Moritz, T. P. Devereaux, and J. K. Freericks, *Phys. Rev. B* **81**, 165112 (2010).
- [18] M. Eckstein, M. Kollar, and P. Werner, *Phys. Rev. B* **81**, 115131 (2010).
- [19] B. Moritz, T. P. Devereaux, and J. K. Freericks, *Comp. Phys. Comm.* **182**, 109 (2011); B. Moritz, A. F. Kemper, M. Sentef, T. P. Devereaux, and J. K. Freericks, preprint (2012), arXiv:1207.3835.

- [20] R. van Leeuwen, N. E. Dahlen, G. Stefanucci, C.-O. Almblad, and U. von Barth, preprint (2005), arXiv:cond-mat/0506130.
- [21] M. Sentef, A. Kemper, B. Moritz, J. Freericks, and T. Devereaux, To be published (2012).
- [22] W. Schülke, G. Stutz, F. Wohlert, and A. Kaprolat, Phys. Rev. B **54**, 14381 (1996).
- [23] B. Moritz, F. Schmitt, W. Meevasana, S. Johnston, E. M. Motoyama, M. Greven, D. H. Lu, C. Kim, R. T. Scalettar, Z.-X. Shen, and T. P. Devereaux, New Journal of Physics **11**, 093020 (2009).